

Magnetic Screening in Hot Non-Abelian Gauge Theory

A. Cucchieri, F. Karsch and P. Petreczky
Fakultät für Physik, Universität Bielefeld,
P.O. Box 100131, D-33501 Bielefeld, Germany

February 1, 2008

Abstract

We analyze the large distance and low-momentum behavior of the magnetic gluon propagator of the SU(2) gauge theory at finite temperature. Lattice calculations within the 4-dimensional as well as the effective, dimensionally reduced 3-dimensional gauge theories in generalized Landau gauges and MAG show that the magnetic propagator is strongly infrared suppressed in Landau gauges but stays large and finite in MAG. Despite these differences in the low-momentum behavior of the propagator calculated in different gauges the magnetic fields are exponentially screened in all gauges considered. From the propagator calculated in maximally Abelian gauge we find for the screening mass, $m_M = (1.48 \pm 0.17)T$ at $T = 2 T_c$.

PACS numbers: 11.10.Wx, 11.15.Ha, 12.38.Mh

Magnetic fields in hot non-abelian gauge theories have to be screened at high temperature for any perturbative description of the plasma phase to make sense [1]. In fact, there is plenty of evidence from analytic [2, 3] as well as numerical [4, 5] calculations for the screening of electric and magnetic fields in non-abelian gauge theories at high temperature. The way this is realized on the partonic level, however, is poorly understood. Our intuitive understanding of screening at high temperature is largely influenced by the perturbative concept of electric and magnetic screening masses [6]. However, while the former is calculable at least to leading order in perturbation theory the latter is intrinsically non-perturbative. Its existence has been postulated to render the perturbative expansion finite [1].

In the vacuum the gluon propagator in Landau gauge is expected to be infrared suppressed. The existing rigorous bounds on the unrenormalized gluon propagator in momentum space imply that it is less singular than p^{-2} in 4 dimension and

p^{-1} in 3 dimension [7] and it has further been argued that it is likely to vanish at zero momentum in the thermodynamic limit [8]. Numerical evidence for this has recently been found in lattice simulations of the $SU(N)$ gauge theory in 3 [9] and in 4 dimensions [10, 11]. As it is well established that the $4d$, $SU(N)$ gauge theory at finite temperature is well described by an effective, $3d$ gauge theory with an adjoint Higgs field [5, 12, 13, 14] and that, moreover, the magnetic sector is little influenced by the scalar Higgs fields [5, 13, 14], the findings for the gluon propagator in 3 dimensions are of direct relevance for the behavior of the magnetic gluon propagator in the $4d$ gauge theory at finite temperature. In fact, it recently has been suggested that also at finite temperature the gluon propagator might be infrared suppressed [15].

It is the purpose of this letter to clarify the long-distance, low-momentum structure of the magnetic gluon propagator of the finite temperature $SU(2)$ gauge theory in $4d$, establish the connection to the $3d$ gauge theory and analyze to what extent these findings are gauge dependent. To this end we will analyze the magnetic gluon propagator in Landau as well as maximally Abelian gauges (MAG). Both gauges are complementary in so far as the bounds on the zero momentum limit of the gluon propagator derived in Landau gauge [7] are not valid in MAG because in this case the Fadeev-Popov matrix turns out to be quadratic in the gauge fields. Although details of the infrared behavior will turn out to be different in different gauges we show that in all cases considered magnetic fields are exponentially screened at large distances. In particular, we will make contact to earlier numerical calculations, which gave evidence for exponential damping of the magnetic propagator at large distances and led to the determination of a magnetic screening mass [4].

We have performed numerical calculations for the $4d$, $SU(2)$ gauge theory at finite temperature as well as for the effective, $3d$ gauge theory with an adjoint Higgs field and the pure $SU(2)$ gauge theory in $3d$. The magnetic gluon propagator, $D_M(z)$, and its Fourier transform, $\bar{D}_M(p)$, have been calculated in Landau gauge and MAG [16] on lattices of size $\Omega = N_\sigma^2 N_z$, in $3d$ and $\Omega = N_\sigma^2 N_z N_\tau$ in $4d$, respectively. In our $3d$ simulations we also have considered an anisotropic generalization of the Landau gauge defined by the gauge condition,

$$\partial_1 A_1 + \partial_2 A_2 + \lambda_3 \partial_3 A_3 = 0 \quad , \quad (1)$$

which for $\lambda_3 = 1$ reduces to the ordinary Landau gauge condition. For the gauge fields we use the straightforward lattice definition, $A_\mu(x) = [U_\mu(x) - U_\mu^\dagger(x)]/2i$, with $U_\mu(x) \in SU(2)$, which reproduces the continuum gauge fields up to $\mathcal{O}(a^2)$ discretization errors. We have checked previously that, up to an overall normalization, other lattice definitions with formally smaller discretization errors lead to identical results [17].

The correlation function for the μ -component of the gauge fields A_μ is defined

in terms of the sum over gauge fields in a hyperplane orthogonal to x_3 ,

$$D_{\mu\mu}(z) = \frac{1}{\Omega} \left\langle \sum_{b=1}^3 \sum_{x_3} Q_\mu^b(x_3 + z) Q_\mu^b(x_3) \right\rangle, \quad (2)$$

with $Q_\mu^b(x_3) = \sum_{x_\perp} \text{Tr}[A_\mu(x_\perp, x_3) \sigma^b]/2$. Here σ^b are the usual Pauli matrices; $x_\perp = (x_1, x_2)$ in $3d$ and $x_\perp = (x_0, x_1, x_2)$ in $4d$, respectively. In terms of this we define the magnetic gluon propagator in coordinate space,

$$D_M(z) = \frac{1}{2} (D_{11}(z) + D_{22}(z)) \quad , \quad (3)$$

and the momentum space propagator

$$\tilde{D}_M(p) = \frac{\beta a^2}{12} \sum_{\mu=1}^3 \sum_{z=0}^{N_z-1} e^{2\pi i k z/N_z} D_{\mu\mu}(z) \quad , \quad (4)$$

with $p = 2 |\sin(\pi k/N_z)|$ and $k = 0, 1, \dots, N_z - 1$. Here β is the coupling appearing in the Euclidean lattice action and a is the lattice spacing.

The $4d$ calculations at finite temperature have been performed in the high temperature phase at twice the critical temperature for the deconfinement transition. All calculations have been performed with the standard Wilson gauge action with gauge coupling $\beta = 2.512$ for $N_\tau = 4$ and $\beta = 2.740$ for $N_\tau = 8$. These values correspond to $T = 2 T_c$ [18]. Corresponding calculations in $3d$ have been performed at $\beta = 8$ on lattices of size $16^2 \times 32$, $24^2 \times 48$, $28^2 \times 56$, $32^2 \times 64$, $48^2 \times 64$ and 64^3 . Moreover, we have performed $3d$ calculations at a smaller coupling, $\beta = 5$, on lattices of size 32^3 up to 96^3 . This choice of parameters for calculations within the reduced theory allows to analyze the behavior of correlation functions in large physical volumes which are 40 times larger than what is feasible within our $4d$ simulations. The gauge coupling of the pure gauge theory was chosen according to $g_3^2(T) = g^2(\mu)T$, where $g^2(\mu)$ is 1-loop running coupling constant of the $4d$ $SU(2)$ gauge theory in the $\overline{MS}$ scheme, $\mu = 18.86T$ and for $\Lambda_{\overline{MS}}$ the numerically determined relation to the deconfinement temperature has been used, *i.e.* $T_c = 1.06\Lambda_{\overline{MS}}$ [4]. This choice of g_3^2 ensures a perfect description of the temperature dependence of the spatial string tension of the $4d$ finite temperature theory in terms of the effective $3d$ theory [19] down to temperatures $T \simeq 2T_c$. Furthermore, we have checked on our smaller lattices that the magnetic propagators calculated in the pure $3d$ $SU(2)$ gauge theory and in the complete $3d$ effective theory, *i.e.* in the presence of an adjoint Higgs field, agree within statistical errors. We therefore will restrict our discussion here to results coming from the pure gauge sector alone. Our $3d$ calculations have been performed at $\beta = 5$ and 8 which correspond to lattice spacings $a = 0.2752/T$ and $0.172/T$, respectively. Varying β in our $3d$ calculations and converting to physical units with the above relation for g_3^2 we have checked that cut-off effects are small.

To verify the cut-off independence of our results in $4d$ the calculations have been performed at two different values of the lattice cut-off, *i.e.* we used lattices of size $N_\sigma^2 N_z N_\tau$ with $N_\tau = 4$ and 8 , $N_z = 2N_\sigma$ and $8 \leq N_\sigma \leq 32$. Keeping the physical volume $VT^3 = 2(N_\sigma/N_\tau)^3$ constant we have verified that $D_M(z)$ is cut-off independent within the statistical errors of our calculation.

The gluon propagator has been calculated on 1000 to 8000 gauge fixed gauge field configurations in the case of $3d$ and about 400 configurations in $4d$ calculations. In Fig. 1 we show the magnetic propagator, $D_M(z)$, calculated in Landau and maximally Abelian gauges on various size lattices. We note that the $4d$ and $3d$ calculations performed on lattices of similar physical size are in good agreement. While the magnetic gluon propagator calculated in MAG shows no significant volume dependence the Landau gauge propagators are strongly volume dependent for $zT \gtrsim 1$. A similarly strong volume dependence we also observe in other generalized Landau gauges, *i.e.* gauges based on the gauge condition given in Eq. 1 with $\lambda_3 \neq 1$. In fact, at large distances the propagators in generalized Landau gauges, including the Coulomb gauge limit, become negative.

The virtue of the effective $3d$ theory is that we can reach here a much larger physical volume and thus can control the volume dependence still apparent in the comparison of $3d$ and $4d$ data shown in Fig. 1. We have analyzed the large distance behavior of $D_M(z)$ in these gauges in our $3d$ calculations on large lattices of size L^3 , $L = 32, 40, 48, 56, 64, 72$ and 96 at $\beta = 5$. This corresponds to physical volumes of size $VT^3 = 683$ up to $VT^3 = 5464$. In all cases we clearly observe negative correlation functions at large distances. For some of these lattices we show the long-distance part of the $D_M(z)$ in Fig. 2 for $\lambda_3 = 1$. Similar results we find for $\lambda_3 \neq 1$. We note that the volume dependence is small on these large lattices. Moreover, the correlation functions become smaller with increasing lattice size. This is in accordance with our expectation that finite size effects are due to zero mode contributions which disappear in the infinite volume limit. On finite lattices, however, they add a volume-dependent, positive constant to the correlation function in coordinate space.

Our analysis thus gives clear evidence that the magnetic gluon correlation function in coordinate space calculated in generalized Landau gauges becomes negative at large distances, *i.e.* for $zT \gtrsim 1.5$. At $T = 0$ such a behavior is, in fact, expected to occur for the gluon propagator as a consequence of the conjecture made by Zwanziger that the Landau gauge gluon propagator vanishes in the infrared limit. In coordinate space it has been suggested that the propagator takes on the form [7]

$$D_M(z) = Ae^{-m_M z} \cos(bm_M z + c) \quad , \quad (5)$$

Although we can at present not quantify the functional form of the correlation function in generalized Landau gauges, our results are consistent with this ansatz up to distances $zT \simeq 3$. To judge whether the correlation function will indeed

continue to oscillate as suggested by Eq. 5 will, however, require a more detailed analysis of the large distance behavior of $D_M(z)$. In general, for $\tilde{D}_M(p)$ to vanish at $p = 0$ in momentum space one should find $\tan(c) = 1/b$ with the above ansatz. This conjecture, $\lim_{p \rightarrow 0} \tilde{D}_M(p) = 0$, may also be valid at non-zero temperature [15].

In contrast to the complicated structure of the Landau gauge propagator the propagator calculated in MAG does show a simple exponential decay at large distances and does not show any significant volume dependence. We could verify that it stays positive at least up to distance $zT = 3.5$. For $zT > 1.5$ simple exponential fits gave $\chi^2/(d.o.f) = 1.7$ and we also found that local masses became independent of z within statistical errors for $zT > 1$. Best fits with the ansatz given in Eq. 5 were thus obtained for $b = 0$. From this we find for the magnetic screening mass of the magnetic propagator calculated in MAG,

$$\frac{m_M}{T} = 1.48 \pm 0.17 \quad \text{at} \quad T = 2T_c \quad . \quad (6)$$

When taking the absolute value of the Landau gauge propagator this is found to be bounded by the exponentially decaying MAG-propagator. This confirms that also in Landau gauge correlations of magnetic fields are exponentially screened.

The fact that the magnetic gluon propagator calculated in Landau gauge becomes negative at large distances translates into a suppression of the momentum space propagator at small momentum. This is shown in Fig. 3 for the Landau gauge propagators, similar results hold for $\lambda_3 \neq 1$. For $p < T$ the momentum space propagator is sensitive to the volume, while for large momenta ($p > T$) it is essentially independent of the lattice size. Moreover, we note that for small momenta the finite volume effects lead to a decrease of $\tilde{D}(p)$ with increasing volume while the volume dependence is already negligible for $p/T \simeq 1$. The occurrence of a maximum in $\tilde{D}(p)$ for $p/T \simeq 1$ and the infrared suppression of the momentum space propagator at $p = 0$ thus is firmly established by our data.

As expected the volume dependence of the magnetic propagator in momentum space is strongest at $p = 0$. On large lattices $\tilde{D}_M(p)$ reaches a maximum at a non-zero but small value of p/T and keeps decreasing at $p = 0$ with increasing lattice size. This might indicate that $\tilde{D}_M(0)$ could actually vanish in the infinite volume limit. The propagator at vanishing momentum reflects the contribution of zero momentum fluctuations of the gauge fields,

$$\tilde{D}_M(0) = \frac{\beta a^2}{12} \Omega \sum_{\mu,b} \langle (\phi_\mu^b)^2 \rangle \quad \text{with} \quad \phi_\mu^b = \frac{1}{\Omega} \sum_x A_\mu^b(x) \quad . \quad (7)$$

In order for it to vanish in the infinite volume limit the zero mode fluctuations $\langle (\phi_\mu^b)^2 \rangle$ thus have to drop faster than Ω^{-1} . To quantify this behavior we have fitted $\tilde{D}_M(0)T^2$ to a simple ansatz that assumes a power-like dependence on the volume, $\tilde{D}_M(0)T^2 = b(VT^3)^{-z} + c$. At present such fits to our 4d and 3d data can, however,

not rule out a non-vanishing value for the constant c . We stress, however, that the value of $\tilde{D}_M(0)$ itself is of little importance for the observed complex structure of the gluon propagator in Landau gauge in coordinate as well as in momentum space and in any case is gauge dependent. Nonetheless, the behavior found here for the finite- T magnetic gluon propagator is consistent with findings of Ref. [9] for the $T = 0$ gluon propagator of the $3d$, $SU(2)$ gauge theory. There it was shown that the magnetic propagator calculated in Landau gauge is less singular than p^{-1} in the infrared and is likely to vanish in $p \rightarrow 0$, $V \rightarrow \infty$ limit.

The momentum dependence of $\tilde{D}_M(p)$ shown in Fig. 3 also makes clear why earlier calculations on medium size lattices led to the determination of a non-zero magnetic mass. The infrared suppression becomes significant only for momenta $p/T \lesssim 0.5$. The existence of a simple *pole mass* becomes unlikely on the basis of our analysis. However, we stress that magnetic fields in coordinate space are, of course, strongly screened. The magnetic gluon propagator calculated in Landau gauge as well as MAG is exponentially damped. The value for the screening mass extracted from the propagator in MAG gauge is in good agreement with earlier in $4d$ [4] and $3d$ [5] lattice calculations in Landau gauge, although the data analysis in these cases was based on a simple pole ansatz for the magnetic mass.

In the course of the calculations reported here we also have reanalyzed the behavior of the electric gluon propagator in different gauges. In that case no suppression in the infrared has been observed and the electric screening mass has been found to be gauge independent within statistical errors. Details of this calculation as well as a more detailed discussion of our results for the magnetic gluon propagator will be presented elsewhere.

Acknowledgements:

The work has been supported by the TMR network ERBFMRX-CT-970122 and by the DFG under grant Ka 1198/4-1. Our calculations have partly been performed at the HLRS in Stuttgart and the $(PC)^2$ in Paderborn.

References

- [1] A.D. Linde, Phys. Lett. **B96** (1980) 289
- [2] G. Alexanian and V.P. Nair, Phys. Lett. **B352** (1995) 435; W. Buchmüller and O. Philipsen, Nucl. Phys. **B443** (1995) 47; F. Eberlein, Phys. Lett. **B439** (1998) 130; Nucl. Phys. **B550** (1999) 303
- [3] D. Karabali and V.P. Nair, Phys. Lett. **B379** (1996) 141; D. Karabali, C. Kim and V.P. Nair, Nucl. Phys. **B524** (1998) 661

- [4] U.M. Heller, F. Karsch and J. Rank, Phys. Lett. **B355** (1995) 511; U.M. Heller, F. Karsch and J. Rank, Phys. Rev. **D57** (1998) 1438
- [5] F. Karsch, M. Oevers and P. Petreczky, Phys. Lett. **B442** (1998) 291
- [6] D.J. Gross, R.D. Pisarski and L.G. Yaffe, Rev. Mod. Phys. **53** (1981) 43
- [7] D. Zwanziger, Nucl. Phys. **B364** (1991) 127
- [8] D. Zwanziger, Nucl. Phys. **B412** (1994) 657
- [9] A. Cucchieri, Phys. Rev. **D60** (1999) 034508
- [10] D.B. Leinweber, J.I. Skullerud, A.G. Williams and C. Parinello, Phys. Rev. **D58** (1998) 0315011
- [11] L. von Smekal, R. Alkofer and A. Hauck, Phys. Rev. Lett. **79** (1997) 3591 ; Ann. Phys. **267** (1998) 1
- [12] P. Lacock, D.E. Miller and T. Reisz , Nucl. Phys. **B369** (1992) 501; L. Kärkkäinen, P. Lacock, B. Peterson and T. Reisz Nucl.Phys. **B395** (1993) 733
- [13] K. Kajantie, M. Laine, K. Rummukainen and M. Shaposhnikov, Nucl. Phys. **B503** (1997) 357
- [14] M. Laine and O. Philipsen, Phys. Lett. **B459** (1999) 259; A. Hart and O. Philipsen, Nucl. Phys. **B572** (2000) 243.
- [15] I. Zahed and D. Zwanziger, Phys. Rev. **D61** (2000) 037501
- [16] K. Amemiya and H. Suganuma, Phys. Rev. **D60** (1999) 114509
- [17] A. Cucchieri, Proceedings of the International Workshop on *Understanding Deconfinement in QCD* (Edt. D. Blaschke, F. Karsch and C.D. Roberts), World Scientific (Singapore 2000)
- [18] J. Fingberg, U.M. Heller and F. Karsch, Nucl. Phys. **B392** (1993) 493
- [19] G.S. Bali et al., Phys. Rev. Lett. **71** (1993) 3059




Figure 1: Volume dependence of the magnetic propagators in coordinate space calculated in Landau (top) and maximally Abelian (bottom) gauges. The propagators were normalized to 1 at $z = 0$. Shown are results from 3d and 4d simulations, which are compared at similar values of the physical volume in units of T^3 . These volumes correspond to lattices of size $16^2 \times 32 \times 8$, $24^2 \times 48 \times 8$ and $16^2 \times 32 \times 4$ in 4d and to $16^2 \times 32$, $24^2 \times 48$, $32^2 \times 64$ and $48^2 \times 64$ in 3d.




Figure 2: The magnetic gluon propagator calculated in generalized Landau gauges with $\lambda_3 = 1$ at $\beta = 5$. Also shown are data from a calculation in MAG on a $48^2 \times 64$ lattice. We only show the propagators at large distances. They were normalized to 1 at $z = 0$.




Figure 3: The magnetic gluon propagator in momentum space calculated in Landau gauge on $3d$ lattices of various sizes at $\beta = 5$.